




**NB Maxolon / Stemetil/ Haloperidol may worsen Parkinson's symptoms and should be avoided.**

### Device Assisted Therapies

**Duodopa Infusion:** \_\_\_\_ - \_\_\_\_ hrs

Morning dose: \_\_\_\_ mls

Continuous Rate: \_\_\_\_ mls/hr

Extra Dose: \_\_\_\_ mls

No. of cassettes /day: \_\_\_\_

Check Stoma/PEG site: \_\_\_\_

**Apomorphine Infusion:** \_\_\_\_ - \_\_\_\_ hrs

Product: Apomine or Movapo

Flow rate: \_\_\_\_ mls/hr (\_\_\_\_ mg/hr)

Bolus: \_\_\_\_ mls (\_\_\_\_ mg) max no. per day

Skin Check: \_\_\_\_

**Deep Brain Stimulation device:**

- Battery or rechargeable device
- Take with you to hospital details of your DBS, including neurologist, manufacturer, device instructions.
- Always check with your DBS team before having an MRI to ensure the procedure is safe for you.
- EEG and ECG Warning: Turn off DBS device before conducting ECG or EEG.
- For surgery Diathermy precautions apply.

If you have any questions, check with the prescribing clinicians, call the manufacturer (Medtronic 1800 510 6735; Abbott: 1800 727 7846; Boston Scientific: 1833 327 8324)